arXiv:1112.2888v1 [math.PR] 13 Dec 2011

Stochastic derivative and heat type PDEs

Constantin Udrişte[*], Virgil Damian[†] and Ionel Ţevy[‡]

Abstract

In this paper we address again the problem of the connection between multitime Brownian sheet and heat type PDEs. The main results include: the volumetric character of the solutions of the forward (backward) diffusion-like PDEs; the forward mean value of a Brownian process as the solution of the forward heat PDE; the backward mean value of a Brownian process as the solution of the backward heat PDE; the multitime stochastic processes with volumetric dependence; the stochastic partial derivative of a stochastic process with respect to a multitime Wiener process; Hermite polynomials stochastic processes; union of Tzitzeica hypersurfaces (constant level sets of multitime stochastic processes with volumetric dependence). The original results permit to extend the complete integrability theory to multitime stochastic differential systems, using path independent curvilinear integrals and volumetric dependence.

1 Introduction

The papers [2], [3] have studied stochastic curvilinear integrals along all sufficiently smooth curves in $\mathbb{R}^2_+$. The most simple situation is those of increasing curves.

Our papers [9], [13], [14] have extended this point of view to stochastic curvilinear integrals in $\mathbb{R}^m_+$ and to completely integrable stochastic differential systems. These research trends and the original results are based on *Itô-Udrişte stochastic calculus rules*

$$dW^a_t\, dW^b_t = \delta^{ab}\, c_\alpha(t) dt^\alpha, dW^a_t\, dt^\alpha = dt^\alpha\, dW^a_t = 0, dt^\alpha\, dt^\beta = 0,$$

where $a, b = \overline{1, d}; \alpha, \beta = \overline{1, m}$, the parameter $t = (t^1, ..., t^m) \in \mathbb{R}^m_+$ means the *multitime*, δ^{ab} is the *Kronecker symbol*, $v = t^1 \cdots t^m$ is the volume of the hyperparallelepiped $\Omega_{0t} \subset R^m_+$, $c_\alpha(t) = \frac{\partial v}{\partial t^\alpha}$ and the tensorial product $\delta^{ab}\, c_\alpha(t)$ represents the *correlation coefficients*. These formulas can be extended to stochastic processes on Riemannian manifolds or on manifolds with a fundamental tensor of type $(2, 0)$ that determines the correlation coefficients. A similar single-time theory can be found in the paper [12].

---

[*] University *Politehnica* of Bucharest, Faculty of Applied Sciences, Deptartment of Mathematics-Informatics I, eMail: udriste@mathem.pub.ro

[†] University *Politehnica* of Bucharest, Faculty of Applied Sciences, Deptartment of Mathematics-Informatics I, eMail: vdamian@mathem.pub.ro

[‡] University *Politehnica* of Bucharest, Faculty of Applied Sciences, Deptartment of Mathematics-Informatics I, eMail: tevy@mathem.pub.ro

Also some ideas have the roots in the basic book [4] containing the single-time theory of Brownian motion, stochastic integral and differential stochastic equations.

The main objective of the paper is the connection between multitime Brownian sheet and heat type PDEs. Section 2 shows that the solutions of the forward diffusion-like PDEs system have a volumetric character and that the forward mean value of a Brownian process is the solution of the forward heat PDE. Section 3 proves that the solutions of the backward diffusion-like PDEs system have a volumetric character and that the backward mean value of a Brownian process is the solution of the backward heat PDE. Section 4 describes the differentiable multitime stochastic processes as sums of series of Hermite polynomials. Section 5 studies unions of Tzitzeica hypersurfaces and their connection to differentiable multitime stochastic processes. Some related deterministic ideas can be found in the papers [5], [10], [11].

2 Forward diffusion-like PDEs system

Let us now consider a real-valued function $f(t,x)$, $t=(t^1,...,t^m)\in\mathbb{R}^m_+$, $x=(x^1,...,x^d)\in\mathbb{R}^d$, with $f(t,0)=0$, which has continuous partial derivatives of the first order with respect to t^α, and of the second order in x^a. If

$$W_t=(W^a_t),\ t=(t^1,...,t^m)\in\mathbb{R}^m_+,\ a=\overline{1,d}$$

is a multitime vectorial Wiener process [9], then the function f defines a stochastic process $f(t,W_t)$, $t\in\mathbb{R}^m_+$. Let us accept (see [1]) that $f(t,W_t)$ is a solution of the *forward diffusion-like PDEs system*

$$\frac{\partial f}{\partial t^\alpha}(t,W_t)-\frac{1}{2}c_\alpha(t)\,\Delta_x f(t,W_t)=0,\ \text{for}\ \alpha=\overline{1,m}, \tag{2.1}$$

where

$$\Delta_x f=\frac{\partial^2 f}{\partial x^a\partial x^b}\,\delta^{ab}.$$

Theorem 2.1 *The solution f of the forward diffusion-like PDEs system (1) depends on the point $(t^1,...,t^m)$ only through the product of components $v=t^1\cdots t^m$, i.e., it is a function of the volume $v=t^1\cdots t^m$ of $\Omega_{0t}\subset\mathbb{R}^m_+$.*

Proof. From (1) it follows

$$c_\beta(t)\,\frac{\partial f}{\partial t^\alpha}(t,W_t)=c_\alpha(t)\,\frac{\partial f}{\partial t^\beta}(t,W_t),\ \alpha\neq\beta.$$

The general solution of this PDEs system is $f(t)=\varphi(t^1\cdots t^m,W_t)$, $t\in\mathbb{R}^m_+$.

If $v=t^1\cdots t^m$ is the volume of $\Omega_{0t}\subset\mathbb{R}^m_+$ and $g(v,x)\overset{def}{=}\varphi\left(t^1\cdots t^m,x\right)$ has continuous partial derivatives of the first order in v and of the second order in x, then from (1) it follows that g satisfies the *forward multitime heat PDE*

$$\frac{\partial g}{\partial v}-\frac{1}{2}\Delta g=0. \tag{2.2}$$

■

Lemma 2.2 (see [1]) *Let*

$$W_t^x = (W_t^a)^x,\ t = (t^1, ..., t^m) \in \mathbb{R}_+^m,\ x \in R^d,\ a = \overline{1, d}$$

be a family of Brownian sheets with $P\{W_t^x = x\} = 1$ *for* $t \in \partial R_+^m$. *If* $f : R^d \to R$ *is a continuous bounded function and* $u(t, x) = E[f(W_t^x)]$ *is the forward mean value, then* $u(t, x)$ *is a unique bounded solution of forward heat type PDEs*

$$\frac{\partial u}{\partial t^\alpha}(t, x) = \frac{1}{2}\, c_\alpha(t)\Delta_x u(t, x),\ t \in \overset{\circ}{R}{}_+^m;$$

$$u(t, x) = f(x),\ t \in \partial R_+^m.$$

Proof. . Let $K(t; x, y)$ be the transition density of an m-parameter R^d- valued Brownian sheet $W^x(t) = W^x(t^1, ..., t^m)$, starting on ∂R_+^m from $x \in R^d$, under P, and moving to $y \in R^d$ at multitime $t = (t^1, ..., t^m) \in R_+^m$, i.e.,

$$K(t; x, y) = \frac{1}{dy} P[W^x(t) \in dy] = \frac{1}{(2\pi\, t^1 \cdots t^m)^{d/2}} \exp\left(\frac{-\|x - y\|^2}{2\, t^1 \cdots t^m}\right).$$

This function is a solution of the forward heat type PDEs, i.e.,

$$\frac{\partial}{\partial t^\alpha} K(t; x, y) = \frac{1}{2} c_\alpha\, \Delta_x K(t; x, y).$$

Taking into account the definition of the mean value $u(t, x)$, the assumptions on the function f and the differentiability of the improper integral function, we find

$$\frac{\partial u}{\partial t^\alpha} = \frac{\partial}{\partial t^\alpha} \int_{R^d} f(y)\, K(t; x, y)\, dy = \int_{R^d} f(y) \frac{\partial}{\partial t^\alpha} K(t; x, y)\, dy$$

$$= \frac{1}{2} c_\alpha \int_{R^d} f(y)\, \Delta_x K(t; x, y)\, dy$$

$$= \frac{1}{2} c_\alpha\, \Delta_x \int_{R^d} f(y)\, K(t; x, y)\, dy = \frac{1}{2} c_\alpha\, \Delta_x u.$$

By definition and by the continuity and boundedness of the function f we easily have

$$\lim_{s \to t, y \to x} u(s, y) = u(t, x) = f(x),$$

for $t \in \partial R_+^m$ and $x \in R^d$. The existence and uniqueness Theorem for a Cauchy problem associated to heat PDE ends the proof. ■

The mean value function u(t, x) is a function of the volume $v = t^1 \cdots t^m$, i.e., $u(t, x) = g(v, x)$.

3 Backward diffusion-like PDEs system

Let us consider again a real-valued function $f(t,x)$ on $\mathbb{R}^m_+ \times \mathbb{R}^d$, with $f(t,0)=0$, which has continuous partial derivatives of the first order with respect to t^α, $\alpha=\overline{1,m}$ and of the second order in x^a. If

$$W_t = (W^a_t),\ \ t=(t^1,...,t^m)\in\mathbb{R}^m_+,\ a=\overline{1,d}$$

is a multitime vectorial Wiener process, then the function f defines a stochastic process

$$y_t = f(t,W_t),\ t\in\mathbb{R}^m_+.$$

By Itô-Udrişte Lemma [9], [13], [14], we can write

$$dy_t = \left(\frac{1}{2}c_\alpha(t)\ \Delta_x f(t,W_t) + \frac{\partial f}{\partial t^\alpha}(t,W_t)\right)dt^\alpha + \frac{\partial f}{\partial x^a}(t,W_t)\,dW^a_t, \tag{3.1}$$

where

$$\Delta_x f = \frac{\partial^2 f}{\partial x^a \partial x^b}\,\delta^{ab}.$$

In order that the stochastic process y_t be a martingale, the drift coefficients in formula (1) must vanish, i.e., f is a solution of the *backward diffusion-like system*

$$\frac{\partial f}{\partial t^\alpha}(t,W_t) + \frac{1}{2}c_\alpha(t)\ \Delta_x f(t,W_t) = 0,\ \text{for}\ \alpha=\overline{1,m}. \tag{3.2}$$

Theorem 3.1 *The solution f of the backward diffusion-like system (4) depends on the point $(t^1,...,t^m)$ only through the product of components $v=t^1\cdots t^m$, i.e., it is a function of the volume $v=t^1\cdots t^m$ of $\Omega_{0t}\subset\mathbb{R}^m_+$.*

Proof. From (4) it follows

$$c_\beta(t)\frac{\partial f}{\partial t^\alpha}(t,W_t) = c_\alpha(t)\frac{\partial f}{\partial t^\beta}(t,W_t),\ \alpha\neq\beta.$$

The general solution of this PDEs system is $f(t)=\varphi(t^1\cdots t^m, W_t),\ t\in\mathbb{R}^m_+$. ■

Let us take the shape of volumetric features into account, recalling that a volumetric function is invariant under the subgroup of central equi-affine (i.e., volume-preserving with no translation) transformations, where the determinant of the representing matrix is 1.

If $v=t^1\cdots t^m$ is the volume of $\Omega_{0t}\subset\mathbb{R}^m_+$ and $g(v,x)\overset{def}{=}\varphi\left(t^1\cdots t^m,x\right)$ has continuous partial derivatives of the first order in v and of the second order in x, then from (3.2) it follows that g satisfies the *backward multitime heat PDE*

$$\frac{\partial g}{\partial v} + \frac{1}{2}\Delta g = 0. \tag{3.3}$$

Consequently, the relation (3.1) reduces to

$$dg(v,W_t) = \frac{\partial g}{\partial x^a}(v,W_t)\,dW^a_t.$$

Thus, if $\mathbb{E}\left[\|\frac{\partial g}{\partial x}(v, W_t)\|^2\right]$ is bounded with respect to t, in bounded subsets of $\mathbb{R}^m_+$, then the stochastic process $g(v, W_t)$ is differentiable with respect to W_t^a and the partial stochastic derivative is $\frac{\partial g}{\partial x^a}(v, W_t)$.

Let T be the maturity multitime of the option.

Lemma 3.2 *Let*

$$W_t^x = (W_t^a)^x,\ t = (t^1, ..., t^m) \in \mathbb{R}^m_+,\ x \in R^d,\ a = \overline{1, d}$$

be a family of Brownian sheets with $P\{W_t^x = x\} = 1$ *for* $t \in \partial R^m_+$. *If* $f : R^d \to R$ *is a continuous bounded function and* $w(t, x) = E_{t,x}[f(W_T^x)]$ *is the backward mean value, then* $w(t, x)$ *is a unique bounded solution of backward heat type PDEs*

$$\frac{\partial w}{\partial t^\alpha}(t, x) = -\frac{1}{2}c_\alpha(t)\Delta_x w(t, x),\ t \in \overset{\circ}{R^m_+};\ w(T, x) = f(x)$$

(terminal value problem).

Proof. Let $V = T^1 \cdots T^m$ and $v = t^1 \cdots t^m$. The transition density is now

$$L(t; x, y) = \frac{1}{(2\pi(V - v))^{d/2}} \exp\left(\frac{-\|x - y\|^2}{2(V - v)}\right).$$

■

Remark 3.3 *The heat PDE has two variants, forward and backward, with the forward one being more common.*

4 Path independent stochastic curvilinear integral

The foregoing theory and the papers [2], [9], [13], [14] suggest to introduce the notion of multitime differentiable stochastic processes. For that we need a multitime vectorial Wiener process

$$W_t = (W_t^a),\ t = (t^1, ..., t^m) \in \mathbb{R}^m_+,\ a = \overline{1, d}.$$

Definition 4.1 *Let* $\gamma_{0t} \subset \mathbb{R}^m_+$ *be an increasing curve joining the points* $0, t \in R^m_+$. *A multitime stochastic process* $\Phi(t) = \Phi(t, W_t),\ t \in \mathbb{R}^m_+$ *is called differentiable with respect to* W_t, *on* $\mathbb{R}^m_+$, *if there exists a multitime vectorial adapted mesurable process* $\phi(t) = (\phi_a(t, W_t)),\ t \in \mathbb{R}^m_+$ *such that* $\mathbb{E}\left[\|\phi(t)\|^2\right]$ *is bounded for* t *in compact sets of* $\mathbb{R}^m_+$ *and*

$$\Phi(t) = \Phi(0) + \int_{\gamma_{0t}} \phi_a(s)\, dW_s^a, \tag{4.1}$$

as a path independent stochastic curvilinear integral.

In terms of stochastic differentials, the multitime stochastic process Φ_t is differentiable if the stochastic system

$$d\Phi(t) = \phi_a(t)\, dW_t^a.$$

is completely integrable, i.e., $\Phi(t)$ is a function of $v = t^1 \cdots t^m$ and hence $\phi(t)$ is a vectorial function of v [2], [13].

The multitime process $\phi_a(t)$ is called the *stochastic partial derivative* of the multitime process $\Phi(t)$ with respect to W_t^a (see also [2], [3]).

Remark 4.2 *The differentiable multitime stochastic processes have properties similar to holomorphic functions: a differentiable process has a differentiable derivative, the curvilinear integral primitive of a differentiable process is differentiable, and each differentiable process admits a power series expansion.*

Theorem 4.3 *Let $w(t,x) = E_{t,x}[f(W_T^x)]$ be the backward mean value. For any Wiener process Y_t, the stochastic process $w(v, Y_t)$ is differentiable.*

Proof. Based on the previous Lemma, the backward mean value function $w(v, Y_t)$ is a solution of the backward multitime heat PDE (terminal value problem)

$$\frac{\partial g}{\partial v} + \frac{1}{2}\Delta g = 0,\ \ g(T,x) = f(x). \tag{4.2}$$

∎

5 Hermite polynomials and stochastic processes

There is a special class of solutions of the backward multitime heat equation which will be particularly interesting in stochastic problems. These are the Hermite polynomials (see also [6], [15]).

To develope further our theory, we need a point $t = (t^1, ..., t^m) \in \mathbb{R}_+^m$, $v = t^1 \cdots t^m$, a point $x = (x^1, ..., x^d) \in \mathbb{R}^d$, a point $n = (n_1, ..., n_d) \in N_0^d$, called multiindex, and the notations $|n| = n_1 + ... + n_d$ and $n! = n_1! \cdots n_d!$. Also for $\xi = (\xi^1, ..., \xi^d) \in R^d$, we write $\xi^n = (\xi^1)^{n_1} ... (\xi^d)^{n_d}$. To simplify, the partial derivative

$$\frac{\partial^{|n|}}{\partial_{\xi^{n_1}} ... \partial_{\xi^{n_d}}} f$$

will be denoted by $\partial_{\xi^n} f$. Now, let us use the generating function expansion

$$e^{<x,\xi> - \frac{1}{2} v \|\xi\|^2} = \sum_{n \in N_0^d} H_n(v,x)\xi^n,\ \ x, \xi \in R^d,\ \ v \in R_+. \tag{5.1}$$

Since

$$H_n(v,x) = \frac{1}{n!}\, \partial_{\xi^n}\, e^{<x,\xi> - \frac{1}{2} v \|\xi\|^2}|_{\xi=0},$$

it follows that $H_n(v,x)$ is a polynomial of degree n in $x = (x^1, ..., x^d)$; it is called the *n-th Hermite polynomial*. On the other hand, the generating function $e^{<x,\xi> - \frac{1}{2} v \|\xi\|^2}$ is a tensor product of d one-dimensional generating functions of the form $e^{x\xi - \frac{1}{2} v \xi^2}$. Consequently

$$H_n(v; x^1, ..., x^d) = H_{n_1}(v; x^1) \cdots H_{n_d}(v; x^d),$$

where H_{n_k} denotes the n_k-th one-dimensional Hermite polynomial.

Completing the square in the definition of generating function, the n^{th} Hermite polynomial of the variables $v \in R_+$ and $x \in R^d$ can be written as

$$H_n(v, x) = \frac{(-v)^n}{n!} e^{\frac{\|x\|^2}{2v}} \partial_{\xi^n} e^{-\frac{1}{2v}\|x - v\xi\|^2}|_{\xi=0} = \frac{(-v)^n}{n!} e^{\frac{\|x\|^2}{2v}} \partial_{x^n} e^{-\frac{\|x\|^2}{2v}}. \tag{5.2}$$

Here and in the rest of the paper, we write $v^n = v^{n_1} \cdots v^{n_d}$.

The sequence $\{H_n(v, \cdot)\}_{n=0}^{\infty}$ is a complete orthogonal system with respect to the weight $(2\pi v)^{\frac{-1}{2}} e^{\frac{-\|x\|^2}{2v}}$. The orthogonality means

$$\mathbb{E}\left[H_m(v, W_t) H_n(v, W_t)\right] = \begin{cases} 0, & \text{if } m \neq n, \\ \dfrac{v^n}{n!}, & \text{if } m = n. \end{cases} \tag{5.3}$$

We introduce the multiindex $k_a = (0, ..., k, ..., 0)$, with k in the place a. Then, using the generating function expansion, taking the partial derivatives with respect to v and x^a, and equating the coefficients, we find

$$\frac{\partial}{\partial v} H_n = -\frac{1}{2} \sum_a H_{n-2_a}, \quad \frac{\partial}{\partial x^a} H_n = H_{n-1_a}.$$

Consequently each Hermite polynomial $H_n(v, x)$ is a solution of the backward multitime heat PDE. Thus

Theorem 5.1 *Each Hermite polynomial $H_n(v, W_t)$ is a differentiable process and its partial derivative with respect to W_t^a is $H_{n-1_a}(v, W_t)$.*

It follows that finite sums of Hermite polynomials processes are differentiable processes. We extend this statement to series of Hermite polynomials, following the ideas of Cairoli and Walsh [2].

Theorem 5.2 *Suppose $\{a_n\}$ is a multiindex family of real numbers such that*

$$\sum_{n \in N_0^d} a_n^2 \frac{v^n}{n!} < \infty, \quad \textit{for all } v > 0 \textit{ (convergent series).}$$

Then, the process Φ_t defined by

$$\Phi_t = \sum_{n \in N_0^d} a_n H_n(v, W_t) \tag{5.4}$$

is differentiable relative to W_t and its partial derivative ϕ_{ta} with respect to W_t^a is given by

$$\phi_{ta} = \sum_{n-1_a \in N_0^d} a_n H_{n-1_a}(v, W_t). \tag{5.5}$$

The convergence of series is understood in L^2.

Proof. By the orthogonality relations (5.3), we find

$$\mathbb{E}\left[\left(\sum_{|n|=0}^{|p|} a_n H_n\left(v, W_t\right)\right)^2\right] = \sum_{|n|=0}^{|p|} a_n^2 \, \frac{v^n}{n!} \, .$$

This mean value is bounded due to the convergence of the series in the right hand member. It follows that the series (5.4) converges in L^2 and the same is true for the series (5.5). Consider now the sequence of partial sums

$$\phi_{ta}^{(|p|)} = \sum_{|n-1_a|=0}^{|p|} a_n H_{n-1_a}\left(v, W_t\right),$$

and let

$$\Phi_t^{(|p|)} = a_0 + \int_{\gamma_{0t}^{\alpha}} \phi_{sa}^{(|p|)} \, dW_s^a = a_0 + \sum_{|n-1_a|=0}^{|p|} a_n H_n\left(v, W_t\right),$$

where

$$\gamma_{0t}^{\alpha} : \, t^1 = c^1, ..., t^{\alpha-1} = c^{\alpha-1}, t^{\alpha} = \tau^{\alpha} \in \left[0, t^{\alpha}\right], t^{\alpha+1} = c^{\alpha+1}, ..., t^m = c^m,$$

is an increasing curve joining the points 0 and t, with $c^{\beta} = const > 0$, $\beta \neq \alpha$. It follows

$$\lim_{|p|\to\infty} \Phi_t^{(|p|)} = \Phi_t \text{ in } L^2.$$

To finish the proof, we need only check that

$$\lim_{|p|\to\infty} \int_{\gamma_{0t}^{\alpha}} \phi_{sa}^{(|p|)} \, dW_s^a = \int_{\gamma_{0t}^{\alpha}} \lim_{|p|\to\infty} \phi_{sa}^{(|p|)} \, dW_s^a, \, \alpha = \overline{1, m}.$$

Again, by the orthogonality relation (5.3), we have

$$\mathbb{E}\left[\left(\phi_{ta} - \phi_{ta}^{(|p|)}\right)^2\right] = \sum_{|n|=|p|+1}^{\infty} a_n^2 \, \frac{v^{n-1_a}}{(n-1_a)!} \, ,$$

and consequently

$$\mathbb{E}\left[\left(\int_{\gamma_{0t}} \left(\phi_{sa} - \phi_{sa}^{(|p|)}\right) dW_s^a\right)^2\right] =$$

$$= \int_0^v \sum_{|n|=|p|+1}^{\infty} a_n^2 \, \frac{v^{n_1} \cdots v^{n_a-1} \cdots v^{n_d}}{(n-1_a)!} \, dv_a = \sum_{|n|=|p|+1}^{\infty} a_n^2 \, \frac{v^n}{n!}.$$

In other words,

$$\lim_{|p|\to\infty} \int_{\gamma_{0t}} \phi_{sa}^{(|p|)} \, dW_s^a = \int_{\gamma_{0t}} \phi_{sa} \, dW_s^a.$$

■

Theorem 5.3 *Suppose that $\varphi(v, x)$ has continuous partial derivatives of the first order in v and of the second order in x, and that $\{\varphi(v, W_t)\}$ is a differentiable process, where $v = t^1 \cdots t^m$. Then, for each $t \in \mathbb{R}^m_+$, we can write*

$$\varphi(v, W_t) = \sum_{|n|=0}^{\infty} a_n H_n(v, W_t),$$

where the convergence is understood in L^2 and, for $t = (t^1, ..., t^m) \in \mathbb{R}^m_+$,

$$a_n = \frac{n!}{v}^{n} \ \mathbb{E}\left[\varphi(v, W_t) H_n(v, W_t)\right]. \tag{5.6}$$

Proof. Define the process $\Phi_t = \sum_{|n|=0}^{\infty} a_n H_n(v, W_t)$. Due to the convergence

$$\sum_{|n|=0}^{\infty} a_n^2 \ \frac{v^n}{n!} < \infty,$$

the series defining Φ_t converges in L^2 and the process Φ_t is differentiable.

Since H_n is an orthogonal sequence, we get

$$\mathbb{E}\left[\Phi_t H_n(v, W_t)\right] = \mathbb{E}\left[a_n H_n^2(v, W_t)\right] = a_n \ \frac{v^n}{n!}.$$

But, by hypothesis,

$$\mathbb{E}\left[\varphi(v, W_t) H_n(v, W_t)\right] = \frac{v^n}{n!} \ a_n.$$

Thus,

$$\mathbb{E}\left[(\varphi(v, W_t) - \Phi_t) H_n(v, W_t)\right] = 0,$$

for all $n \in \mathbb{N}_{\not{k}}$. Then, $\varphi(v, W_t) - \Phi_t \equiv 0$. ■

6 Union of Tzitzeica hypersurfaces

A hypersurface $M \subset \mathbb{R}^m_+$, $m \geq 3$, is called *Tzitzeica hypersurface* [7], [8], provided there exists a constant $a \in \mathbb{R}$ such that we have $K = a\, d^{m+1}$, for all points $t = (t^1, ..., t^m) \in M$, where K is the Gauss curvature of the hypersurface and d is the distance from the origin of the space to the tangent hyperplane to the hypersurface at the current point t. Since the Gauss curvature K describes the shape of the hypersurface, a Tzitzeica hypersurface has a bending against the tangent hyperplane in fixed proportion to the normal component of the position vector t. The most simple Tzitzeica hypersurfaces are the constant level sets $M_c : t^1 \cdots t^m = c$ in $\mathbb{R}^m$ (2^{m-1} connected components).

Remark 6.1 *The Gauss curvature of a Cartesian implicit surface*

$$M_c : F(t^1, t^2, t^3) = c$$

in $\mathbb{R}^3$ is

$$K = \Big[[F_3(F_3F_{11} - 2F_1F_{13}) + F_1^2F_{33}][F_3(F_3F_{22} - 2F_2F_{23}) + F_2^2F_{33}$$

$$-\,(F_3(-F_1F_{23}+F_3F_{12}-F_yF_{13})+F_1F_2F_{33})^2\Big]\,[F_3^2(F_1^2+F_2^2+F_3^2)^2]^{-1}.$$

The surface M_c is curving like a paraboloid if $K(t) > 0$, hyperboloid if $K(t) < 0$, or a cylinder or plane if $K(t) = 0$, near a point $t = (t^1, t^2, t^3) \in M_c$.

Let us show that the constant level sets of the functions $\varphi(t^1 \cdots t^m, x)$, with respect to $t = (t^1, ..., t^m) \in \mathbb{R}_+^m$, are Tzitzeica hypersurfaces or unions of simple Tzitzeica hypersurfaces in $\mathbb{R}_+^m$ indexed by the points x.

Theorem 6.2 *If $\varphi : \mathbb{R} \to \mathbb{R}$ is a C^2 nonconstant function and c is a noncritical value of φ, then the constant level set $N_c : \varphi(t^1 \cdots t^m) = c$ is a union of simple Tzitzeica hypersurfaces.*

Proof. . Let A_c the set of the solutions of the equation $\varphi(v) = c$ and $v = t^1 \cdots t^m$. Then the constant level set N_c is the union of the constant level sets $t^1 \cdots t^m = k,\ k \in A_c$. If c is not a critical value of φ, then the set N_c is a union of hypersurfaces. ■

Remark 6.3 *If c is a critical value of φ, then the set N_c is a union of constant level sets $t^1 \cdots t^m = k,\ k \in A_c$, but those level sets which correspond to $\varphi'(k) = 0$ are not hypersurfaces.*

Theorem 6.4 *Each section $x = c$ of the hypersurface $H_n(v, x) = 0$ in $\mathbb{R}^{m+d} = \{(t, x)\}$ is a union of Tzitzeica hypersurfaces in $\mathbb{R}_+^m$.*

The constant level set

$$\mathbb{E}\,[H_n\,(v, W_t)\,H_n\,(v, W_t)] = c > 0$$

is a simple Tzitzeica hypersurface.

References

[1] H. Allouba, *From Brownian time Brownian sheet to a fourth order and a Kuramoto-Sivashinski-variant interacting PDEs systems*. arXiv 1007 5302v2 [math. PR] 26 Apr. 2011, 1-19.

[2] R. Cairoli, J. Walsh, *Stochastic integrals in the plane*. Acta Math., **134** (1975), 111–183.

[3] M. Dozzi, *Stochastic processes with a multidimensional parameter*. Longman Scientific & Technical, 1989.

[4] G. da Prato, *Introduction to differential stochastic equations*. Scuola Normale Superiore, 1998.

[5] L. Matei, C. Udrişte, *Multitime sine-Gordon solitons via geometric characteristics*. *Balkan Journal of Geometry and Its Applications*, **16** (2011), 81-89.

[6] S. Stojanovic, *Computational Financial Mathematics using Mathematica*. Birkhäuser, 2003.

[7] G. Tzitzeica, *Sur une nouvelle classe de surfaces*. C. R. Acad. Sc., **150** (1910), 955-956.

[8] G. Tzitzeica, *Sur une nouvelle classe de surfaces*. C. R. Acad. Sc., **150** (1910), 1227-1229.

[9] C. Udrişte, *Multitime Stochastic Control Theory*. Selected Topics on Circuits, Systems, Electronics, Control & Signal Processing. Edited by S. Kartalopoulos, M. Demiralp, N. Mastorakis, R. Soni, H. Nassar. WSEAS Press (2007), 171-176.

[10] C. Udrişte, V. Arsinte, C. Cipu *Von Neumann analysis of linearized discrete Tzitzeica PDE*. Balkan Journal of Geometry and Its Applications, **15** (2010), 100-112.

[11] C. Udrişte, V. Arsinte C. Cipu, *Tzitzeica and sine-Gordon solitons*. in Balkan Journal of Geometry and Its Applications, **16** (2011), 150-154.

[12] C. Udrişte, V. Damian, *Simplified single-time stochastic maximum principle*. Balkan Journal of Geometry and Its Applications, **16** (2011), 155-173.

[13] C. Udrişte, V. Damian, *Multitime stochastic maximum principle based on curvilinear integral actions*. Manuscript, 2011.

[14] C. Udrişte, V. Damian, L. Matei, I. Ţevy, *Multitime differentiable stochastic processes, diffusion PDE, Tzitzeica hypersurfaces*. U.P.B. Sci. Bull., Series A, *to appear*.

[15] G. Zitkovic, *Topics in Stochastic Analysis, Lecture 1: Hermite Polynomials*. Available at: http://www.ma.utexas.edu/users/gordanz/teaching/lecture1.pdf (2011)